

# Homology cylinders in knot theory

Hiroshi GODA

(Tokyo University of Agriculture and Technology)

and


Takuya SAKASAI

(Tokyo Institute of Technology)


December 19, 2009

# Introduction (1)

## §1. Introduction

- $\Sigma_{g,1} =$    $(g \geq 0, \text{ oriented})$

with a standard cell decomposition:



## Definition (Goussarov, Habiro, Garoufalidis-Levine, Levine)


$(M, i_+, i_-)$  : a *homology cylinder* (HC) over  $\Sigma_{g,1}$

$\stackrel{\text{def}}{\iff} \left\{ \begin{array}{l} M : \text{a compact oriented 3-manifold,} \\ i_+, i_- : \Sigma_{g,1} \hookrightarrow \partial M \text{ two embeddings (markings)} \end{array} \right.$

satisfying

- 1  $i_+$ : orientation-preserving,  $i_-$ : orientation-reversing;
- 2  $\partial M = i_+(\Sigma_{g,1}) \cup i_-(\Sigma_{g,1})$ ,  
 $i_+(\Sigma_{g,1}) \cap i_-(\Sigma_{g,1}) = i_+(\partial\Sigma_{g,1}) = i_-(\partial\Sigma_{g,1})$ ;
- 3  $i_+|_{\partial\Sigma_{g,1}} = i_-|_{\partial\Sigma_{g,1}}$ ;
- 4  $i_+, i_- : H_*(\Sigma_{g,1}; \mathbb{Z}) \xrightarrow{\cong} H_*(M; \mathbb{Z})$  isomorphisms.

- $(M, i_+, i_-)$  : a homology cylinder (over  $\Sigma_{g,1}$ )



## Definition

$$\mathcal{C}_{g,1} := \{(M, i_+, i_-) : \text{HC over } \Sigma_{g,1}\} / (\text{marking pres. diffeo}).$$

## Stacking


For  $(M, i_+, i_-), (N, j_+, j_-) \in \mathcal{C}_{g,1}$ ,

$$(M, i_+, i_-) \cdot (N, j_+, j_-) := (M \cup_{i_- \circ (j_+)^{-1}} N, i_+, j_-) \in \mathcal{C}_{g,1}$$

$\leadsto \mathcal{C}_{g,1}$  becomes a **monoid**.

unit:  $(\Sigma_{g,1} \times [0, 1], \text{id} \times 1, \text{id} \times 0)$

where corners of  $\Sigma_{g,1} \times [0, 1]$  are rounded, and



## Examples

①  $\mathcal{M}_{g,1}$ : the mapping class group of  $\Sigma_{g,1}$

$[\varphi] \in \mathcal{M}_{g,1}$ , i.e.

$\varphi : \Sigma_{g,1} \xrightarrow{\sim} \Sigma_{g,1}$ : a diffeo. s.t.  $\varphi|_{\partial\Sigma_{g,1}} = \text{id}$

$\implies (\Sigma_{g,1} \times I, \text{id} \times 1, \varphi \times 0) \in \mathcal{C}_{g,1}$ .

We can check

$\mathcal{M}_{g,1} \hookrightarrow \mathcal{C}_{g,1}$  : monoid embedding

$\rightsquigarrow \mathcal{C}_{g,1}$  is an *enlargement* of  $\mathcal{M}_{g,1}$ .

- 2 surgery along **clovers** (Goussarov) or **claspers** (Habiro)
- 3 surgery along **pure string links** (Habegger, Levine)
- 4 connected sum with a homology 3-sphere  $X$ :  
 $(\Sigma_{g,1} \times [0, 1] \# X, \text{id} \times 1, \text{id} \times 0) \in \mathcal{C}_{g,1}$ .

Today we focus on


- 5 complementary sutured manifolds of Seifert surfaces of **a special class of knots**.

# Homological fibered knots (1)

## §2. Homological fibered knots


- $K \subset S^3$  : a knot,  
 $S$  : a Seifert surface of  $K$  of genus  $g$ .

$M_S$  : the cobordism obtained from  $E(K)$  by cutting along  $S$   
= the **(complementary) sutured manifold** for  $S$ .





- By using any identification  $i : \Sigma_{g,1} \xrightarrow{\cong} S$ , we obtain a *marked* sutured manifold  $(M_S, i_+, i_-)$ :



Question When this becomes a homology cylinder?

## Proposition (Crowell-Trotter, ..., Goda-S.)

$K$  : a knot in  $S^3$ ,

$K$  has a Seifert surface  $S$  of genus  $g$  s.t.  $M_S$  is a homology product (over a copy of  $S$ )

$\iff$  The following hold:

- $S$  is a minimal genus Seifert surface,
- The Alexander polynomial  $\Delta_K(t)$  of  $K$  is monic,
- $\deg(\Delta_K(t)) = 2 \text{ genus}(K)$ .

## Definition

A knot  $K$  in  $S^3$  is said to be *homologically fibered* if

- (1)  $\Delta_K(t)$  is monic,
- (2)  $\deg(\Delta_K(t)) = 2 \text{ genus}(K)$ .

## Remarks

- (Fibered knots)  $\subset$  (HFknots [Homological Fibered knots]) corresponds to  $\mathcal{M}_{g,1} \subset \mathcal{C}_{g,1}$ .
- We can define *rational* homological fibered knots ( $\mathbb{Q}$ -HFknot) by assuming only (2).

## “Uniqueness”

### Proposition

$K$  : an HFknot of genus  $g$

$S_1, S_2$  : minimal genus Seifert surfaces


For any markings of  $\partial M_{S_1}$  and  $\partial M_{S_2}$ ,  $\exists N \in \mathcal{C}_{g,1}$  s.t.

$$M_{S_1} \cdot N = N \cdot M_{S_2} \in \mathcal{C}_{g,1}$$

In particular, any monoid homomorphism

$$\mathcal{C}_{g,1} \rightarrow A \quad \text{w/ } A: \text{ an abelian group}$$

gives an invariant of HFknots.



Pretzel knot  $P(-3, 5, 9)$  is an HFknot.

Easy to see  $P(-2n + 1, 2n + 1, 2n^2 + 1)$  is an HFknot.

## §3. Factorization formulas of Alexander invariants

### Classical case

- $K \subset S^3$  : a knot,  
   $S$  : a Seifert surface of  $K$  w/ a Seifert matrix  $A$ .

Assume that  $A$  is invertible over  $\mathbb{Q}$  (i.e.  $K$  is a  $\mathbb{Q}$ -HFknot).

Then

$$\begin{aligned}\Delta_K(t) &= \det(A^T - tA) \\ &= \det(A^T) \det(I_{2g(S)} - t(A^T)^{-1}A)\end{aligned}$$

What does this *factorization* mean?

We can check:

- $A^T$  and  $A$  represent

$$i_+, i_- : \mathbb{Z}^{2g} \cong H_1(\Sigma_{g,1}) \longrightarrow H_1(M_S) \cong \mathbb{Z}^{2g}$$

under certain bases of  $H_1(\Sigma_{g,1})$  and  $H_1(M_S)$ . In fact,

$$\begin{aligned} \det(A) &= \text{The top (bottom) coeff. of } \Delta_K(t) \\ &= \pm |H_1(M, i_+(\Sigma_{g,1}))| \\ &= \tau(C_*(M_S, i_+(\Sigma_{g,1}); \mathbb{Q}) \quad \text{torsion} \end{aligned}$$

- $\sigma(M_S) := (A^T)^{-1}A \in Sp(2g, \mathbb{Q})$ .  
(Can regard  $\sigma(M_S)$  as an  $H_1$ -monodromy of  $M_S$ .)

So, roughly speaking, our factorization formula says

$$\begin{aligned}\Delta_K(t) &= \det(A^T) \det(I_{2g(S)} - t(A^T)^{-1}A) \\ &= (\text{torsion of } M_S) \cdot (\text{effect of } H_1\text{-monodromy of } M_S).\end{aligned}$$

Remark By Milnor,

$$\frac{\Delta_K(t)}{1-t} = \tau_{\mathbb{Z}}(K),$$

where  $\tau_{\mathbb{Z}}(K)$  is the **Reidemeister torsion** associated with the  $\mathbb{Z}$ -cover of  $E(K)$ .



- For an HFknot  $K$ ,

$$\begin{aligned}\Delta_K(t) &= \det(A^T) \det(I_{2g(S)} - t(A^T)^{-1}A) \\ &= \pm \det(I_{2g(S)} - t(A^T)^{-1}A).\end{aligned}$$

~> The factorization formula is useless for HFknots!

~> We will give a generalization by using twisted homology.

## Higher-order case (Twisted coefficients)

- $K$ : an HFknot,
- $M_S = (M_S, i_+, i_-) \in \mathcal{C}_{g,1}$  : an HC associated with  $K$ ,
- $\mathcal{K} := \text{Frac}(\mathbb{Z}H_1(M_S)) \cong \mathbb{Q}(t_1, \dots, t_{2g})$  as twisted coefficients.

### Lemma

For  $\pm \in \{+, -\}$ ,  $H_*(M_S, i_\pm(\Sigma_{g,1}); \mathcal{K}) = 0$ .

cf. classical case:  $H_*(M_S, i_\pm(\Sigma_{g,1}); \mathbb{Z}) = 0$ .

## Definition

- The  **$\mathcal{K}$ -torsion**  $\tau_{\mathcal{K}}(M_S)$  is

$$\tau_{\mathcal{K}}(M_S) := \tau(C_*(M_S, i_+(\Sigma_{g,1}); \mathcal{K})) \in GL(\mathcal{K}) / \sim.$$

- The **Magnus matrix**  $r_{\mathcal{K}}(M_S) \in GL(2g, \mathcal{K})$  is the representation matrix of the right  $\mathcal{K}$ -isom.:

$$\begin{array}{ccc}
 H_1(\Sigma_{g,1}, \mathcal{K}) & \xrightarrow[i_-]{\cong} & H_1(M_S, \mathcal{K}) & \xrightarrow[i_+^{-1}]{\cong} & H_1(\Sigma_{g,1}, \mathcal{K}) \\
 \parallel & & & & \parallel \\
 \mathcal{K}^{2g} & \xrightarrow[r_{\mathcal{K}}(M_S)]{\cong} & & & \mathcal{K}^{2g}
 \end{array}$$

Remark By substituting  $t_i \mapsto 1$ , we have

$$\tau_{\mathcal{K}}(M_S) \mapsto \det A = \pm 1, \quad r_{\mathcal{K}}(M_S) \mapsto \sigma(M_S).$$

- If  $K$  is fibered w/ the monodromy  $\varphi \in \mathcal{M}_{g,1}$ , then

$$r_{\mathcal{K}}(M_S) = \overline{\left( \frac{\partial \varphi(\gamma_j)}{\partial \gamma_i} \right)_{1 \leq i, j \leq 2g}}.$$

## Theorem (Fibered obstructions)

$K, M_S$ : as before.

If  $K$  is **fibered**, then

- 1 all the entries of the Magnus matrix  $r_{\mathcal{K}}(M_S)$  are **Laurent polynomials** in  $\mathbb{Q}[t_1^{\pm}, \dots, t_{2g}^{\pm}] \subset \mathcal{K} = \mathbb{Q}(t_1, \dots, t_{2g})$ ,
- 2 the  $\mathcal{K}$ -torsion  $\tau_{\mathcal{K}}(M_S)$  is **trivial**.

## Theorem (Factorization formula)

Let

$$\rho : \pi_1(E(K)) \longrightarrow \frac{\pi_1(E(K))}{\pi_1(E(K))''} \cong H_1(M_S) \rtimes H_1(E(K))$$

be the natural projection, and let  $t \in H_1(E(K))$  be a meridian loop. Then we have

$$\begin{aligned} \tau_{\mathcal{K}(t^\pm; \sigma)}(E(K)) &= \frac{\tau_{\mathcal{K}}(M_S) \cdot (I_{2g} - t \cdot r_{\mathcal{K}}(M_S))}{1 - t} \\ &\in \mathcal{K}_1(\mathcal{K}(t^\pm; \sigma)) / \pm \rho(\pi_1(E(K))), \end{aligned}$$

where LHS is the **noncommutative higher-order torsion** associated with  $\rho$  (defined by Cochran, Harvey and Friedl).

## §4. 12-crossings non-fibered homological fibered knots

### Facts on fibered knots vs. HFknots

- HFknots with at most 11-crossings are all fibered.
- There are 13 non-fibered HFknots with 12-crossings. In particular, Friedl-Kim showed that these 13 knots are not fibered by using twisted Alexander polynomial associated with finite representations.

We computed  $r_{\mathcal{K}}(M_S)$  and  $\tau_{\mathcal{K}}(M_S)$  for these 13 knots and checked that **each of them also detects the non-fiberedness of all 13 HFknots.**

## Recipe

- 1 Get all the pictures of those 13 knots.  
[By Computer (Database (KnotInfo) on Internet)]
- 2 For each of them,
  - 1 Find a minimal genus Seifert surface  $S$ .  
[By hand]
  - 2 Calculate an **admissible** presentation of  $\pi_1(M_S)$ .  
[By hand]
  - 3 Compute  $r_{\mathcal{K}}(M_S)$  and  $\tau_{\mathcal{K}}(M_S)$ .  
[By hand and also by computer program]


# 12-crossings non-fibered homological fibered knots (3)

## 1 List of non-fibered HFknots:


Knot 12n-	Genus	Alexander polynomial
0057	2	$1 - 2t + 3t^2 - 2t^3 + t^4$
0210	3	$1 - t - t^2 + 3t^3 - t^4 - t^5 + t^6$
0214	3	$1 - t - t^2 + 3t^3 - t^4 - t^5 + t^6$
0258	2	$1 - 4t + 5t^2 - 4t^3 + t^4$
0279	2	$1 - 6t + 11t^2 - 6t^3 + t^4$
0382	2	$1 - 5t + 7t^2 - 5t^3 + t^4$
0394	2	$1 - 6t + 11t^2 - 6t^3 + t^4$
0464	2	$1 - 4t + 5t^2 - 4t^3 + t^4$
0483	2	$1 - 4t + 5t^2 - 4t^3 + t^4$
0535	2	$1 - 7t + 11t^2 - 7t^3 + t^4$
0650	2	$1 - 4t + 7t^2 - 4t^3 + t^4$
0801	2	$1 - 5t + 7t^2 - 5t^3 + t^4$
0815	2	$1 - 2t + t^2 - 2t^3 + t^4$




- 2 Example of calculation of admissible presentation



12n0057





# 12-crossings non-fibered homological fibered knots (5)





Generators	$i_-(\gamma_1), \dots, i_-(\gamma_4), z_1, \dots, z_{10}, i_+(\gamma_1), \dots, i_+(\gamma_4)$
Relations	$z_1 z_5 z_6^{-1}, z_2 z_3 z_4 z_1, z_3 z_9^{-1} z_5^{-1}, z_7 z_4 z_8^{-1}, z_8 z_{10} z_6,$ $z_2 z_5 z_7^{-1} z_5^{-1}, z_9 z_4 z_{10}^{-1} z_4^{-1}, i_-(\gamma_1) z_1^{-1} z_5^{-1}, i_-(\gamma_2) z_2,$ $i_-(\gamma_3) z_4 z_8 z_7 z_5^{-1}, i_-(\gamma_4) z_4, i_+(\gamma_1) z_5^{-1}, i_+(\gamma_2) z_9^{-1} z_6^{-1},$ $i_+(\gamma_3) z_6 z_4 z_7 z_5^{-1} z_3^{-1} z_5 z_6^{-1}, i_+(\gamma_4) z_6 z_7^{-1} z_6^{-1}$

# 12-crossings non-fibered homological fibered knots (6)





12n0210





12n0214

# 12-crossings non-fibered homological fibered knots (7)




12n0258





12n0279


# 12-crossings non-fibered homological fibered knots (8)




12n0382





12n0394




# 12-crossings non-fibered homological fibered knots (9)




12n0464




12n0483



# 12-crossings non-fibered homological fibered knots (10)





12n0535



12n0650

# 12-crossings non-fibered homological fibered knots (11)





## Computational results for 12n0057

$$r_{\mathcal{K}}(M_S) = \begin{pmatrix} \frac{x^3+x^1 x^2^2 (-1+x^2 (-1+x^4)) - x^2 x^3 x^4}{x^1 x^2^2 (-1+x^2 (-1+x^4))} & - \frac{(-1+x^4) (-1+x^2 x^4)}{-1+x^2 (-1+x^4)} & \frac{x^4}{1+x^2-x^2 x^4} & 0 \\ - \frac{(1+x^1 x^2) x^3}{x^1^2 x^2 (-1+x^2 (-1+x^4))} & - \frac{x^2 (1+x^1 x^2) (-1+x^4)}{x^1 (-1+x^2 (-1+x^4))} & - \frac{(1+x^2) (1+x^1 x^2^2 (-1+x^4))}{x^1 x^2 (-1+x^2 (-1+x^4))} & \frac{1}{x^4} \\ \frac{x^3}{x^1 (-1+x^2 (-1+x^4))} & \frac{x^2^2 (-1+x^4)}{-1+x^2 (-1+x^4)} & \frac{x^2 (1+x^2) (-1+x^4)}{-1+x^2 (-1+x^4)} & 0 \\ \frac{(x^1 x^2^2 - x^3) x^4}{x^1^2 x^2 (-1+x^2 (-1+x^4))} & \frac{x^2 x^4 (x^1 x^2 + x^3 - x^3 x^4)}{x^1 x^3 (-1+x^2 (-1+x^4))} & \frac{(1+x^2) (x^1 x^2^2 - x^3) x^4}{x^1 x^2 x^3 (-1+x^2 (-1+x^4))} & 1 \end{pmatrix},$$

$$\tau_{\mathcal{K}}(M_S) = x^1 x^2^4 + x^1 x^2^5 - x^1 x^2^5 x^4,$$

where  $x^j = i_+(\gamma_j)$ .

Each of  $r_{\mathcal{K}}(M_S)$  and  $\tau_{\mathcal{K}}(M_S)$  shows that 12n0057 is not fibered!

## §5. Further results and projects

- Detection of non-fiberedness by Johnson-Morita homomorphism:  
(2nd Johnson homomorphism + *Yokomizo cokernel*)
- Local moves among HFknots???
- *Categorification* of factorization formulas???

$$\begin{array}{ccc} \widehat{HFK}(K) & \rightsquigarrow & \Delta_K(t), \\ & \begin{array}{c} ? \\ \rightsquigarrow \end{array} & \tau_{\mathcal{K}(t^{\pm}; \sigma)}(E(K)), \\ SFH(M_S, K) & \rightsquigarrow & \tau_{\mathcal{K}}(M_S), \\ ??? & \rightsquigarrow & r_{\mathcal{K}}(M_S). \end{array}$$

*decategorification*

Fin